

PSEUDOKARST - K.G. Grimes

tropical climates (e.g. Wall & Wilford, 1966; Dragovich, 1968); in Tasmania a cave produced by solution of secondary minerals in an alteration zone of a dolerite, described by Hale & Spry (1964); volcano-karst, a form produced when labile minerals in newly erupted volcanic ashes are rapidly dissolved to produce weird sculpturing forms (Fairbridge, 1968 b, and Bourke in this volume); and laterite-karst, where silicate minerals are removed in solution from within a deep weathering profile (Grimes, 1974). (Laterite-karst is distinct from the laterite caves in West Australia described by Lefroy & Lake (1972) which appear to be a piping phenomena).

MORPHOLOGICAL KARST

produced by




Table 1. Classification of Karst and Pseudokarst

(B) True pseudokarst

The different types of pseudokarst are grouped here according to the phase in which the material is removed (see Table 1):

(i) *Solid phase removal.* The greatest variety of pseudokarsts result from the removal of solid material. Piping is a form of soil erosion which can also occur in poorly consolidated deposits. The process is one in which clay particles are removed in suspension by groundwater percolating through cracks or pore spaces in the sediment. In conjunction with this headward sapping occurs at springs. Once underground tunnels are produced by these processes they can be enlarged by the normal corrosive action of streams flowing through them. Fletcher et al (1954) postulated five conditions for the development of piping: (1) a source of water, (2) a surface infiltration rate which exceeds the permeability rate of some subsoil layer, (3) the existence of an erodable layer just above the retarding layer, (4) the water above the retarding layer must have a hydraulic gradient along which it can flow laterally, and (5) there must be an outlet for the lateral flow. The process is facilitated by the presence of swelling clay minerals and by a high exchangeable sodium potential. Piping can result in intricate forms of morphological karst, including caves, underground streams, dolines, etc. (Parker et al, 1964; Parker, 1968; Mears, 1968; Feininger, 1969). The process is characterised by its rapidity when compared with solution karst; Parker et al (op cit) quote rates of 2,700 cu.ft. per year for enlargement of Officer's Cave, U.S.A.. Some authors (eg. Khobzi, 1972, and the Cave Summary sheets produced by the A.S.F.) use "suffosion" as a synonym for piping. This is not valid as suffosion is a periglacial phenomenon (see Fairbridge, 1968 c).

Some Australasian examples of piping caves will be described by Bourke and by Shannon elsewhere in this volume. A good local example is the Flagstone Creek Caves (Gillieson, 1971).

Sea Caves result from the combined hydraulic action of waves and the pressure of compressed air, aided by salt spray weathering. Dolines can form above the caves. A large number of sea caves in N.S.W. have been described in recent years and Toomer & Welch will be summarising these studies elsewhere in this symposium.

The mechanical removal of material from animal burrows or by human mining activities could be considered as a form of pseudokarst. Mine tunnels are analagous in form to natural caves and disused mines are often occupied by bats and other cave dwelling species. They can develop speleothem deposits if conditions are suitable and these may include some of uncommon mineralogy in the case of metalliferous mines. Some beautiful stalactites of malachite and azurite have formed in disused parts of the Mt. Isa mines. Subsidence areas and dolines can form above old mine workings.

Some other forms are fault fissures resulting from tectonic movements, and the formation of rock shelters and small caves by weathering or wind erosion. Weathering can combine with piping of the resultant clay material to enlarge joints in granite and produce an underground drainage, e.g. the Labertouche Cave in Victoria (Ollier, 1965) and the Underground River at Wyberba, Qld., (Pound, 1971).

(ii) *Liquid phase removal.* The most common form in this group is the formation of lava tunnels by the draining of molten lava from within a solidifying flow. Ollier & Brown (1965) have described in some detail the process which formed the Victorian lava caves. In these the lava solidified in laminae separated by still molten material which then became segregated into discrete cylindrical tubes. These were at first filled with molten rock and capable of remelting their walls and moving uphill for short distances under hydrostatic pressure. If the tube were then drained a lava cave would result and sagging or collapse of the roof could form surface depressions. Hatheway (1971) presents a similar

PSEUDOKARST - K.G. Grimes

theory but considers that the mobile cylinders of liquid lava originate at the toe of the flow and develop backwards up the flow to its source. Greeley (1971, 1972) has observed the formation of tunnels by the crusting over of lava channels on the surface of an active flow in Hawaii. In Queensland the best known lava tunnels are in the Mt. Suprise area (Watt, 1972; White, 1965, pp. 91-101). The lava cave at Bunya Mountains may be a weathering feature rather than a true lava tunnel (Graham, 1971).

Melting of ice and snow produces several types of pseudokarst. Meltwater caves in snow banks or in glaciers are of most interest to speleologists. Glacier caves can reach 2.5 km length (Halliday & Anderson, 1970). Local examples have been described from the Snowy Mountains by Halbert & Halbert (1972) and from New Zealand (Shannon, 1972). An interesting form is the summit firn cave of Mt. Rainier, U.S.A., in which the melting is mainly due to heat from volcanic activity (Kiver & Mumma, 1971).

In arctic areas seasonal thawing of frozen ground results in a terrain known as thermokarst (Dylik, 1968; Anderson & Hussey, 1963) which includes pits, dry gullies, small hummocks and closed depressions. The processes involved include subsurface mudflows which erupt at the surface as mud volcanoes and this is the preferred use of the term "suffosion" (Fairbridge, 1968 c).

(iii) *Gaseous removal*. This is a theoretical possibility, as in the ablation of snow and ice, but I don't know of any examples where this is the dominant process. Ablation could have formed the steam cups and fluting of the ice walls of the caves on Mt. Rainier (Kiver & Mumma, op cit).

References

The following list will provide an introduction to the various types of pseudokarst and non-limestone karst. Perusal of Australian Speleological Abstracts will disclose a large number of Australian pseudokarst examples.

- ANDERSON, G.S., & HUSSEY, K.M., 1963 Preliminary investigation of thermokarst development on the North Slope, Alaska. *Proc. Iowa Acad.Sci.*, 70, 306-20.
- DRAGOVICH, D., 1968 Granite lapies at Remarkable Rocks, South Australia. *Rev. Geomorphol.Dyn.*, 18(1), 8-16.
- DYLIK, J., 1968 Thermokarst. In *The Encyclopedia of Geomorphology*. Ed. R.W. Fairbridge, New York, Reinhold. Pp.1149-51.
- FAIRBRIDGE, R.W., 1968 a Salt Karst. In *The Encyclopedia of Geomorphology*. Ed. R.W. Fairbridge, New York, Reinhold. Pp.967-8.
- FAIRBRIDGE, R.W., 1968 b Volcano-karst. In *The Encyclopedia of Geomorphology*. Ed. R.W. Fairbridge, New York, Reinhold. Pp.1205-8.
- FAIRBRIDGE, R.W., 1968 c Suffosion and Tundra Craters. In *The Encyclopedia of Geomorphology*. Ed. R.W. Fairbridge, New York, Reinhold. Pp.1099-1100.
- FEININGER, T., 1969 Pseudokarst on quartz diorite, Colombia. *Z.Geomorph.*, 13(3), 287-96.
- FLETCHER, J.E., HARRIS, K., PETERSON, M.B., & CHANDLER, V.N., 1954 Piping. *Trans.Am.Geophys.Union*, 35(2), 258-62.
- GILLIESON, D.S., 1971 Flagstone Creek Caves. *Down Under*, 10(4), 93-4.
- GRAHAM, A., 1971 A preliminary note on a cave in basalt, Bunya Mountains National Park, Queensland. *Helictite*, 9(4), 73-6.
- GREELEY, R., 1971 Observations of actively forming lava tubes and associated structures, Hawaii. *Modern Geol.*, 2, 207-33.
- GREELEY, R., 1972 Additional observations of actively forming lava tubes and associated structures, Hawaii. *Modern Geol.*, 3, 157-60.
- GRIMES, K.G., 1974 The Mesozoic and Cainozoic geology of the Lawn Hill, Westmoreland, Mornington, and Cape Van Dieman sheet areas. *Bur. Miner. Resour. Aust. Rec.*, 1974/106 (unpubl.)
- HATHEWAY, A.W., 1971 Lava tubes and Collapse Depressions. (Ph.D. Thesis, Univ. Arizona). *Diss.Abstr.Int.*, 32(4), 2234B. (Abstr.)

PSEUDOKARST - K.G. Grimes

- HALBERT, E., & HALBERT, S., 1972 Meltwater caves in the Snowy Mountains. *J.Syd.Speleo.Soc.*, 16(8), 223-4.
- HALE, G.E., & SPRY, A., 1964 A cave in dolerite at Wayatinah, Tasmania. *J.Geol.Soc.Aust.*, 11(2) 213-6.
- HALLIDAY, W.R. & ANDERSON, C.H., 1970 Glacier Caves - a new field of Speleology. *Stud.Speleol.*, 2(2), (extract printed in *Oolite*, 2(3), 12-13)
- KHOBZI, J., 1972 Erosion chimique et mecanique dans la genese de depressions "pseudo-karstiques" souvent endoreiques. *Revue Geomorph.Dyn.*, 21(2), 57-70.
- KIVER, E.P., & MUMMA, M.D., 1971 Summit Firn Caves, Mount Rainier, Washington. *Science*, 173, 320-2.
- LEFROY, T., & LAKE, P., 1972 A laterite cave system in the Chittering Region, West Australia. *Western Caver*, 12(3), 68-77.
- MEARS, B., 1968 Piping. In *The Encyclopedia of Geomorphology*. Ed. R.W. Fairbridge, New York, Reinhold. Pp.849-50.
- MONROE, W.H., 1970 A glossary of Karst Terminology. *U.S. Geol.Surv. Water Supply Pap.*, 1899K.
- OLLIER, C.D., 1965 Some features of granite weathering in Australia. *Z.Geomorphol.*, 9, 285-304.
- OLLIER, C.D., & BROWN, M.C., 1965 Lava caves of Victoria. *Bull.Vulcanol.*, 28, 215-29.
- PARKER, G.G., 1968 Officer's Cave, Eastern Oregon, Revisited. *Geol.Soc.Am.Spec.Pap* 101, p415 (Abstr.).
- PARKER, G.G., SHOWN, L.M., & RATZLOFF, K.W., 1964 Officer's Cave, a pseudo-karst feature in altered tuff and volcanic ash of the John Day Formation in Eastern Oregon. *Geol.Soc.Am.Bull.*, 75, 393-402.
- POUND, M., 1971 Wyberba (trip report). *Down Under*, 10(3), 67-8.
- SHANNON, C.H.C., 1972 Meltwater caves in the glaciers of the Southern Alps. *Down Under*, 11(2), 36-8.
- SWEETING, M.M., 1972 Karst Landforms. London, Macmillan.
- WALL, J.R.D., & WILFORD, G.E., 1966 A comparison of small-scale solution features on microgranodiorite and limestone in West Sarawak, Malaysia. *Z.Geomorph.*, 10, 462-8.
- WATT, A., 1972 Lava Caves in North Queensland, Einasleigh-Mt.Suprise area. *Proc. 8th Biennial Conf.Aust.Speleo.Fedn.*, 18-24.
- WHITE, D.A., 1965 Geology of the Georgetown/Clark River area, Queensland. *Bur.Miner.Resour.Aust.Bull.*, 71, 91-101.

Additional references not mentioned in text:

- GIBSON, I.L., 1974 Blister Caves associated with an Ethiopian Volcanic Ash-Flow Tuff. *Stud.Speleol.*, 2(6), 225-232.
- HALLIDAY, W.R., & ANDERSON, C.H., 1970 Glacier Caves, a new field of Speleology. *Stud.Speleol.*, 2(2), 53-59.
- PETERSON, D.W., & SWANSON, D.A., 1974 Observed Formation of Lava Tubes during 1970-71 at Kilauea Volcano, Hawaii. *Stud.Speleol.*, 2(6), 209-222.